



## The use of light-emitting diode therapy in the treatment of photoaged skin.

Baez F<sup>1</sup>, Reilly LR.

### Author information

### Abstract

**BACKGROUND:** Light-emitting diode (LED) therapy is an increasingly popular methodology for the treatment of sun damage. Combination use of light wavelengths reported to stimulate collagen synthesis and accelerate fibroblast-myofibroblast transformation may display a composite rejuvenative effect.

**OBJECTIVE:** To clinically assess reduction in sun damage signs following a 5-week course of LED therapy and to assess subject's perception of the treatment.

**METHODS:** Thirteen subjects with wrinkles or fine lines in the periorbital and nasolabial region and those presenting Glogau scale photodamage grade II-III received nine 20-min duration light treatments using the Omnilux LED system. The treatments combined wavelengths of 633 and 830 nm at fluences of 126 and 66 J/cm<sup>2</sup>, respectively. Sun-damage reduction was assessed at 6, 9, and 12 weeks by clinical photography and patient satisfaction scores.

**RESULTS:** The majority of subjects displayed "moderate" (50%) or "slight" (25%) response to treatment at investigator assessment. Treatment of the periorbital region was reported more effective than the nasolabial region. At 12-week follow-up, 91% of subjects reported improved skin tone, and 82% reported enhanced smoothness of skin in the treatment area.

**CONCLUSION:** Good response to LED therapy has been shown in this modest sample. Larger trials are needed to assess optimum frequency of light treatments and overall treatment time.

PMID: 17760698 DOI: [10.1111/j.1473-2165.2007.00329.x](https://doi.org/10.1111/j.1473-2165.2007.00329.x)

[Indexed for MEDLINE]

Publication type, MeSH terms

LinkOut - more resources

0 comments

[How to join PubMed Commons](#)